

ECZEMA FLARE: CARE PLAN

< 1 year of age

The NZ Child and Youth Eczema Clinical Network | Te Rōpū Kiripai Hapori

Place patient sticker here

FOR MORE INFO:



Date: _____

Visit
kidshealth.org.nz
for eczema
videos

BATH:

- Once a day. No soap or bubble bath.
- Use soap-free wash or moisturiser as a soap substitute.
- Dilute bleach bathing, if prescribed: ☐

MOISTURISER:

- Use every day, all over.
- Apply morning and night and with every nappy change.
- Apply when needed to soothe itch.
- Can be used instead of soap.

YOUR MOISTURISER IS:

Prescriber to tick

- ☐ Cetomacrogol & Glycerol (Sorbolene) ☐ Fatty cream
- ☐ Paraffin 50:50 ☐ Cetomacrogol
- ☐ Other: _____

TOPICAL STEROID: FACE

- Once a day to eczema that is red/darker and itchy.
- Do not use Aristocort on eyelids.
- Use plain Vaseline for irritation around the mouth.

YOUR FACE STEROID IS:

Prescriber to tick

- ☐ 1% hydrocortisone ☐ Aristocort
- ☐ Other: _____

TOPICAL STEROID: BODY, ARMS & LEGS

- Once a day to ALL eczema that is rough, red/darker and itchy.
- You can mix half and half with moisturiser to help it spread over large areas.
- Don't use under the nappy unless told to by your doctor.

YOUR BODY STEROID IS:

Prescriber to tick

- ☐ 1% hydrocortisone ☐ Aristocort
- ☐ Advantan
- ☐ Other: _____

TREATMENT PLAN:

- Apply steroids on days with ☒
- Prescriber to circle start day.
- See your prescriber for review in 3 - 4 weeks to make a long term management plan.

	MON	TUES	WEDS	THURS	FRI	SAT	SUN
WEEK 1	☒	☒	☒	☒	☒	☒	☒
WEEK 2	☒	☒	☒	☒	☒	☒	☒
WEEK 3	☐	☒	☐	☒	☐	☒	☐
WEEK 4	☒	☐	☒	☐	☒	☐	☒
WEEKS 5+	☐	☐	☐	☐	☐	☒	☒

From week 5 onwards: EITHER use steroid to eczema as needed at the first sign of a flare OR if eczema patches flare often, use steroid as a preventer 2 days per week (Sat and Sun)



ECZEMA FLARE: CARE PLAN

< 1 year of age

The NZ Child and Youth Eczema Clinical Network | Te Rōpū Kiripai Hapori

MOISTURISER

Cetomacrogol & Glycerol (Sorbolene)



Fatty Cream



Paraffin 50:50



Cetomacrogol



STERIODS

FACE OR BODY

1% Hydrocortisone



MILD

Aristocort
(triamcinolone acetonide 0.02%)



MODERATE

BODY

Locoid
(hydrocortisone butyrate 0.1%)




MODERATE

Advantan
(methylprednisolone aceponate 0.1%)



STRONG

Beta
(betamethasone valerate 0.1%)



STRONG

Elocon
(mometasone furoate 0.1%)



STRONG